This article was downloaded by: ["University at Buffalo, the State University of New York (SUNY)"] On: 26 October 2011, At: 07:16 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Synthetic Communications

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lsyc20

# Aluminum(III) Hydrogensulfate: An Efficient Solid Acid Catalyst for the Preparation of 5-Substituted 1H-Tetrazoles

S. Mohammad Sajadi<sup>a b</sup>, Mehdi Khalaj<sup>c</sup>, Sayyed Mohammad Hosseini Jamkarani<sup>d</sup>, Mehdi Mahame<sup>e</sup> & Mehrdad Kashefib<sup>b</sup> <sup>a</sup> Department of PetroleumgeoSciences, Faculty of Sciences, Soran University, Soran, Kurdistan Regional Government, Irag

<sup>b</sup> Department of Biology, Faculty of Sciences, Soran University, Soran, Kurdistan Regional Government, Irag

<sup>c</sup> Department of Chemistry, Islamic Azad University, Buinzahra Branch, Qazvin, Iran

<sup>d</sup> Department of Chemistry, Islamic Azad University, Karaj Branch, Karaj, Iran

<sup>e</sup> Department of Chemistry, Islamic Azad University, Aliabad Katool Branch, Aliabad Katool, Iran

Available online: 11 Jul 2011

To cite this article: S. Mohammad Sajadi, Mehdi Khalaj, Sayyed Mohammad Hosseini Jamkarani, Mehdi Mahame & Mehrdad Kashefib (2011): Aluminum(III) Hydrogensulfate: An Efficient Solid Acid Catalyst for the Preparation of 5-Substituted 1H-Tetrazoles, Synthetic Communications, 41:20, 3053-3059

To link to this article: <u>http://dx.doi.org/10.1080/00397911.2010.516861</u>

# PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-conditions</u>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



Synthetic Communications<sup>®</sup>, 41: 3053–3059, 2011 Copyright © Taylor & Francis Group, LLC ISSN: 0039-7911 print/1532-2432 online DOI: 10.1080/00397911.2010.516861

## ALUMINUM(III) HYDROGENSULFATE: AN EFFICIENT SOLID ACID CATALYST FOR THE PREPARATION OF 5-SUBSTITUTED 1*H*-TETRAZOLES

# S. Mohammad Sajadi,<sup>1,2</sup> Mehdi Khalaj,<sup>3</sup> Sayyed Mohammad Hosseini Jamkarani,<sup>4</sup> Mehdi Mahame,<sup>5</sup> and Mehrdad Kashefib<sup>2</sup>

 <sup>1</sup>Department of PetroleumgeoSciences, Faculty of Sciences, Soran University, Soran, Kurdistan Regional Government, Iraq
<sup>2</sup>Department of Biology, Faculty of Sciences, Soran University, Soran, Kurdistan Regional Government, Iraq
<sup>3</sup>Department of Chemistry, Islamic Azad University, Buinzahra Branch, Qazvin, Iran
<sup>4</sup>Department of Chemistry, Islamic Azad University, Karaj Branch, Karaj, Iran
<sup>5</sup>Department of Chemistry, Islamic Azad University, Aliabad Katool Branch,

Aliabad Katool, Iran

## **GRAPHICAL ABSTRACT**



**Abstract** An efficient method for preparation of 5-substituted 1H-tetrazoles via [2+3] cycloaddition of nitriles and sodium azide is reported using aluminum(III) hydrogensulfate as an effective solid acid. This method has the advantages of good yields, simple methodology, and easy workup.

**Keywords** Aluminum(III); [2+3] cycloaddition; hydrogensulfate; nitrile; solid acid catalyst; 5-substituted 1*H*-tetrazole

#### INTRODUCTION

The growth of tetrazole chemistry over the past 25 years has been significant, mainly as a result of the roles played by tetrazoles in coordination chemistry as ligands, in medicinal chemistry as stable surrogates for carboxylic acids, and in material applications, including explosives, agriculture, and photography.<sup>[1–4]</sup>

Received July 6, 2010.

Address correspondence to S. Mohammad Sajadi, Department of PetroleumgeoSciences, Faculty of Sciences, Soran University, PO Box 862, Soran, Kurdistan Regional Government, Iraq. E-mail: smohammad.sajadi@gmail.com



Scheme 1. Conversion of nitriles to the corresponding 5-substituted 1H-tetrazoles using Al(HSO<sub>4</sub>)<sub>3</sub>.

Another important application of tetrazoles is preparation of imidoylazides.<sup>[5]</sup> The conventional method of synthesizing tetrazoles is by addition of azide ions to organic nitriles or cyanamides.<sup>[6]</sup> Earlier reported methods for the synthesis of 5-substituted tetrazoles suffer from drawbacks such as the use of strong Lewis acids or expensive and toxic metals, and the in situ–generated hydrazoic acid, which is highly toxic and explosive.<sup>[7]</sup> Several syntheses of 5-substituted tetrazoles have been reported through the [2+3] cycloaddition of nitriles using NaN<sub>3</sub> or TMSN<sub>3</sub> in the presence of catalysts such as FeCl<sub>3</sub>–SiO<sub>2</sub>,<sup>[6f]</sup> AlCl<sub>3</sub>,<sup>[8]</sup> (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>O · BF<sub>3</sub>,<sup>[9]</sup> tetrabutyl ammonium fluoride (TBAF),<sup>[10]</sup> Pd(PPh<sub>3</sub>)<sub>4</sub>,<sup>[11]</sup> Zn/Al hydrotalcite,<sup>[12]</sup> and ZnO.<sup>[13]</sup>

The use of solid acid salt catalysts,  $[^{[4]}$  such as Al(HSO<sub>4</sub>)<sub>3</sub>, for synthesizing organic intermediates and fine chemicals is gaining increasing awareness and is a field of intense research activity.  $^{[15]}$  This salt is a stable and nonhygroscopic solid material that is insoluble in most organic solvents.

We herein report a new protocol for preparation of 5-substituted 1*H*-tetrazoles derivatives from nitriles using aluminum(III) hydrogensulfate as a solid acid catalyst (Scheme 1).

#### EXPERIMENTAL

#### General

All reagents were purchased from Merck and Aldrich and used without further purification. <sup>13</sup>C NMR and <sup>1</sup>H NMR spectra were recorded on Brucker 100- and 250-MHz instruments using tetramethylsilane (TMS) as an internal standard. Chemical shifts are reported in parts per million (ppm), and coupling constants are reported in hertz (Hz). Infrared (IR) spectra were recorded on a Shimadzu 470 spectrophot-ometer. Thin-layer chromatography (TLC) was performed on Merck precoated silica-gel 60-F254 plates. Al(HSO<sub>4</sub>)<sub>3</sub> was prepared from AlCl<sub>3</sub> and H<sub>2</sub>SO<sub>4</sub>, according to the literature.<sup>[14c,15]</sup> Massah and coworkers studied the characterization of the Al(HSO<sub>4</sub>)<sub>3</sub> using DSC, TG-DTG, and FT–IR spectroscopy.<sup>[15b]</sup>

## General Procedure for Preparation of 5-Substituted 1H-Tetrazoles

Al(HSO<sub>4</sub>)<sub>3</sub>(10 mol%) was added to a mixture of nitrile (2 mmol), NaN<sub>3</sub> (3 mmol), and distilled dimethylformamide (6 mL) and stirred at 120 °C for the appropriate time (Table 1). After completion of the reaction (as monitored by TLC), the mixture was cooled to room temperature. The solid acid was filtered, and the filtrate was treated with ethyl acetate (35 mL) and 5 N HCl (20 mL) and stirred vigorously. The resultant organic layer was separated, and the aqueous layer was again extracted with ethyl acetate (25 mL). The combined organic layer was washed with water (8 mL)

Entry	Substrate	Product	Time (h)	Yield (%) <sup><i>a</i></sup>	Ref.
1	CN Ia		18	91	6f
2	CN Ib MeO		18	87	6f
3		Me	18	85	6f
4			18	90	6f
5	CI-CN Ie		18	89	6f
6	CN		18	90	6f
7	La CN	H N Br	18	84	6f
8	CN Me		22	85	17
9			36	60	18
10	CI IJ CN		18	84	6f

Table 1. Synthesis of various 5-substituted 1*H*-tetrazoles in the presence of Al(HSO<sub>4</sub>)<sub>3</sub> by reaction of sodium azide and nitriles at 120 °C

(Continued)

Entry	Substrate	Product	Time (h)	Yield $(\%)^a$	Ref.
11	CN		18	85	6f
12			30	69	17

Table 1. Continued

<sup>*a*</sup>Yields refer to the pure isolated products.

and concentrated to give a crude product. Column chromatography using silica gel gave pure product in excellent yield. The pure products were characterized by IR and NMR. All the products are known compounds, and the spectral data and melting points were identical to those reported in the literature.

#### **RESULTS AND DISCUSSION**

The general synthetic method is depicted in Scheme 1. 5-Substituted 1*H*-tetrazoles were obtained from the reaction of nitrile with sodium azide in the presence of  $Al(HSO_4)_3$  as a solid acid catalyst at 120 °C for appropriate time in good yields, as summarized in Table 1.

First, we optimized the amount of Al(HSO<sub>4</sub>)<sub>3</sub> catalyst required in the reaction between benzonitrile and sodium azide (Table 2). Water was not a suitable solvent for this reaction. Not many organic solvents are stable at the high temperatures necessary for cycloaddition reactions (sometimes as high as 130 °C), and for this reason dimethylformamide (DMF) is most commonly used for this purpose.<sup>[2a,9,12]</sup> The optimum amount of Al(HSO<sub>4</sub>)<sub>3</sub> was found to be 10 mol% in the presence of nitrile (2 mmol) and sodium azide (3 mmol) in DMF (6 mL). We next examined a variety of structurally divergent benzonitriles possessing a wide range of functional groups to understand the scope and generality of the Al(HSO<sub>4</sub>)<sub>3</sub>-promoted [2 + 3] cycloaddition reaction to form 5-substituted 1*H*-tetrazoles, and the results are summarized in Table 1. The nature of the substituent on the benzonitrile did not affect the reaction time (Table 1, entries 1–7, 10, and 11). Reactions of the 3-methylbenzonitrile (Table 1, entry 8) and 4-hydroxybenzonitrile (Table 1, entry 9) were complete at 120 °C after 22 and 36 h, respectively, and gave the corresponding tetrazole in excellent yields.

The products were characterized by IR, <sup>1</sup>H NMR, and <sup>13</sup>C NMR spectroscopy and from melting points. The disappearance of one strong and sharp absorption band (CN stretching band) and the appearance of an NH stretching band in the IR spectra were evidence for the formation of 5-substituted 1*H*-tetrazoles. <sup>13</sup>C NMR spectra display signals for tetrazole ring carbons of arylaminotetrazoles in the range of 154–157 ppm (depending on the nature of the substituents in the amino functionality).<sup>[16]</sup>

$Al(HSO_4)_3 \pmod{6}$	Solvent	Yield% <sup>b</sup>
5	DMF	54
7	DMF	76
10	DMF	93
10	DMSO	91
15	DMF	94
0	DMF	$0^c$
	AI(HSO <sub>4</sub> ) <sub>3</sub> (mol%) 5 7 10 10 15 0	Al(HSO <sub>4</sub> ) <sub>3</sub> (mol%)     Solvent       5     DMF       7     DMF       10     DMF       10     DMSO       15     DMF       0     DMF

**Table 2.** Preparation of 5-phenyltetrazole using varying amounts of  $Al(HSO_4)_3$  under thermal conditions at  $120 \degree C^a$ 

<sup>*a*</sup>Reaction conditions: nitrile (2 mmol), NaN<sub>3</sub> (3 mmol), Al(HSO<sub>4</sub>)<sub>3</sub> (10 mol%), and DMF, (6 mL), reaction time (18 h) at 120 °C.

<sup>b</sup>Isolated yield.

<sup>c</sup>In the absence of catalyst at 120 °C, no reaction occurred after 18 h.

#### CONCLUSION

In conclusion, we have developed a novel and highly efficient method for the synthesis of various 5-substituted 1H-tetrazoles by treatment of nitriles with sodium azide in the presence of Al(HSO<sub>4</sub>)<sub>3</sub> as an effective catalyst. The significant advantages of this methodology are good yields, elimination of dangerous and harmful hydrazoic acid, simple workup procedure, and easy preparation and handling of the catalyst.

#### ACKNOWLEDGMENT

The authors acknowledge the financial support by the Soran University.

#### REFERENCES

- Bulter, R. N. In *Comprehensive Heterocyclic Chemistry II*; A. R. Katritzky, C. W. Ress, E. F. V. Scriven, (Eds.); Pergamon: New York, 1996; Vol. 4, p. 621.
- (a) Herr, R. 5-Substituted-1*H*-tetrazoles as carboxylic acid isosteres: Medicinal chemistry and synthetic methods. J. Bioorg. Med. Chem. 2002, 10, 3379–3393; (b) Holland, G. F.; Pereira, J. N. Heterocyclic tetrazoles, a new class of lipolysis inhibitors. J. Med. Chem. 1967, 10, 149–154; (c) Figdor, S. K.; Schach von Wittenau, M. Metabolism of 5-(3-pyridyl)tetrazole. J. Med. Chem. 1967, 10, 1158–1159; (d) Rhonnstad, P.; Wensbo, D. On the relative strength of the 1*H*- tetrazol-5-yl- and the 2-(triphenylmethyl)-2*H*tetrazol-5-yl-group in directed ortho-lithiation. Tetrahedron Lett. 2002, 43, 3137–3139.
- Klapötke, T. M.; Stierstorfer, J.; Weber, B. New energetic material: Synthesis and characterization of copper 5-nitriminotetrazolates. *Inorg. Chim. Acta.* 2009, 362, 2311–2320.
- John, E. O.; Kirchmeier, R. L.; Shreeve, J. M. Reactions of 5-(perfluoro-alkyl)tetrazolates with Cyanogen, Nitrosyl, and Cyanuric Chlorides. *Inorg. Chem.* 1989, 28, 4629.
- Modarresi-Alam, A. R.; Khamooshi, F.; Rostamizadeh, M.; Keykha, H.; Nasrollahzadeh, M.; Bijanzadeh, H. R.; Kleinpeter, E. Dynamic <sup>1</sup>H NMR spectroscopic study of the restricted S-N rotation in aryl-N-(arylsulfonyl)-N- (triphenylphosphoranylidene)imidocarbamates. J. Mol. Struc. 2007, 841, 61–66.
- (a) Kadaba, P. K. Role of protic and dipolar aprotic solvents in heterocyclic syntheses via 1,3-dipolar cycloaddition reactions. *Synthesis* 1973, 71–84; (b) Wittenberger, S. J. Recent developments in tetrazole chemistry: A review. *Org Prep. Proc. Int.* 1994, 26, 499–532;

(c) Curran, D. P.; Hadida, S.; Kim, S. Y. *Tris*(2-perfluorohexylethyl)tin azide: A new reagent for preparation of 5-substituted tetrazoles from nitriles with purification by fluorous/organic liquid–liquid extraction. *Tetrahedron* **1999**, *55*, 8997–9006; (d) Huff, B. E.; Staszak, M. A. A new method for the preparation of tetrazoles from nitriles using trimethylsilylazide/trimethylaluminum. *Tetrahedron Lett.* **1993**, *34*, 8011–8014; (e) Modarresi-Alam, A. R.; Nasrollahzadeh, M. Synthesis of 5-arylamino- 1*H*(2*H*)-tetrazoles and 5-amino-1-aryl-1*H*-tetrazoles from secondary arylcyanamides in glacial acetic acid: A simple and efficient method. *Turk. J. hem.* **2009**, *33*, 1–14; (f) Nasrollahzadeh, M.; Bayat, Y.; Habibi, D.; Moshaee, S. FeCl<sub>3</sub>–SiO<sub>2</sub> as a reusable heterogeneous catalyst for the synthesis of 5-substituted 1*H*-tetrazoles via [2 + 3] cycloaddition of nitriles and sodium azide. *Tetrahedron Lett.* **2009**, *50*, 4435–4438; (g) Nasrollahzadeh, M.; Habibi, D.; Shahkarami, Z.; Bayat, Y. A general synthetic method for the formation of arylaminotetrazoles using natural natrolite zeolite as a new and reusable heterogeneous catalyst. *Tetrahedron* **2009**, *65*, 10715–10719; (h) Habibi, D.; Nasrollahzadeh, M.; Faraji, A. R.; Bayat, Y. Efficient synthesis of arylaminotetrazoles in water. *Tetrahedron* **2010**, *66*, 3866–3870.

- (a) Duncia, J. V.; Pierce, M. E.; Santella III, J. B. Three synthetic routes to a sterically hindered tetrazole: A new one-step mild conversion of an amide into a tetrazole. J. Org. Chem. 1991, 56, 2395–2400; (b) Sisido, K.; Nabika, K.; Isida, T.; Kozima, S. Formation of organotin-nitrogen bonds, III: N-Trialkyltin-5-substituted tetrazoles. J. Organomet. Chem. 1971, 33, 337–346; (c) Carini, D. J.; Duncia, J. V.; Aldrich, P. E.; Chui, A. T.; Johnson, A. L.; Pierce, M. E.; Price, W. A.; Santella III, J. B.; Wells, G. J.; Wexler, R. R.; Wong, P. C.; Yoo, S.-E.; Timmermans, P. B. M. W. M. Nonpeptide angiotensin II receptor antagonists: The discovery of a series of N-(biphenylylmethyl)imidazoles as potent, orally active antihypertensives. J. Med. Chem. 1991, 34, 2525–2547; (d) Wittenberger, S. J.; Donner, B. G. Dialkyltin oxide-mediated addition of trimethylsilyl azide to nitriles: A novel preparation of 5-substituted tetrazols. J. Org. Chem. 1993, 58, 4139–4144.
- Matthews, D. P.; Green, J. E.; Shuker, A. J. Parallel synthesis of alkyl tetrazole derivatives using solid support chemistry. J. Comb. Chem. 2000, 2, 19–23.
- 9. Kumar, A.; Narayanan, R.; Shechter, H. Rearrangement reactions of (hydroxyphenyl)carbenes. J. Org. Chem. 1996, 61, 4462–4465.
- Amantini, D.; Beleggia, R.; Fringuelli, F.; Pizzo, F.; Vaccoro, L. TBAF-catalyzed synthesis of 5-substituted 1*H*-tetrazoles under solventless conditions. *J. Org. Chem.* 2004, 69, 2896–2898.
- Gyoung, Y. S.; Shim, J.-G.; Yamamoto, Y. Regiospecific synthesis of 2-allylated-5-substituted tetrazoles via palladium-catalyzed reaction of nitriles, trimethylsilyl azide, and allyl acetates. *Tetrahedron Lett.* 2000, *41*, 4193–4196.
- Kantam, M. L.; Shiva Kumar, K. B.; Phani Raja, K. An efficient synthesis of 5-substituted 1*H*-tetrazoles using Zn/Al hydrotalcite catalyst. *J. Mol. Catal. A: Chem.* 2006, 247, 186–188.
- Kantam, M. L.; Shiva Kumar, K. B.; Sridhar, C. Nanocrystalline ZnO as an efficient heterogeneous catalyst for the synthesis of 5-substituted 1*H*-tetrazoles. *Adv. Synth. Catal.* 2005, 347, 1212–1214.
- (a) Modarresi-Alam, A. R.; Nasrollahzadeh, M.; Khamooshi, F. Solvent-free preparation of primary carbamates using silica sulfuric acid as an efficient reagent. *Arkivoc* 2007, *16*, 238–245; (b) Modarresi-Alam, A. R.; Khamooshi, F.; Nasrollahzadeh, M.; Amirazizi, H. A. Silica-supported perchloric acid (HClO<sub>4</sub>–SiO<sub>2</sub>): An efficient reagent for the preparation of primary carbamates under solvent-free conditions. *Tetrahedron* 2007, *63*, 8723–8726; (c) Modarresi-Alam, A. R.; Nasrollahzadeh, M.; Khamooshi, F. Al(HSO<sub>4</sub>)<sub>3</sub> as an efficient reagent for the preparation of primary carbamates under solvent-free conditions. *Sci. Iranica* 2008, *15*, 452–455.

#### 5-SUBSTITUTED 1H-TETRAZOLE

- 15. (a) Ghorbani-Choghamarani, A.; Zeinivand, J. Catalytic oxidation of sulfides to sulfoxides using aluminium hydrogen sulfate, sodium nitrite, and catalytic amounts of metal bromide. J. Iran. Chem. Soc. 2010, 7, 190–194; (b) Massah, A. R.; Asadi, B.; Hoseinpour, M.; Molseghi, A.; Javad Kalbasi, R.; Javaherian Naghash, H. A novel and efficient solvent-free and heterogeneous method for the synthesis of primary, secondary, and bis-N-acylsulfonamides using metal hydrogen sulfate catalysts. Tetrahedron 2010, 65, 7696–7705.
- Goljer, I.; Svetlik, J.; Hrusovsky, I. <sup>13</sup>C-NMR of substituted tetrazoles. *Monatsh. Chem.* 1983, 114, 65–70.
- 17. Mihina, J. S.; Herbst, R. M. The reaction of nitriles with hydrazoic acid: Synthesis of monosubstituted tetrazoles. J. Org. Chem. 1950, 15, 1082–1092.
- Demko, P. Z.; Sharpless, K. B. Preparation of 5-substituted 1*H*-tetrazoles from nitriles in water. J. Org. Chem. 2001, 66, 7945–7950.

Downloaded by ["University at Buffalo, the State University of New York (SUNY)"] at 07:16 26 October 2011